arXiv:0908.1851v2 [math.AG] 28 Mar 2010

# HOMOGENEOUS TORIC VARIETIES

IVAN V. ARZHANTSEV AND SERGEY A. GAIFULLIN

Abstract. A description of transitive actions of a semisimple algebraic group $G$ on toric varieties is obtained. Every toric variety admitting such an action lies between a product of punctured affine spaces and a product of projective spaces. The result is based on the Cox realization of a toric variety as a quotient space of an open subset of a vector space $V$ by a quasitorus action and on investigation of the $G$-module structure of $V$.

## 1. Introduction

We study toric varieties $X$ equipped with a transitive action of a connected semisimple algebraic group $G$. In this case $X$ is called a *homogeneous toric variety*. The ground field $\mathbb{K}$ is algebraically closed and of characteristic zero.

Consider a quasiaffine variety

$$\mathcal{X} = \mathcal{X}(n_1, \ldots, n_m) := (\mathbb{K}^{n_1} \setminus \{0\}) \times \cdots \times (\mathbb{K}^{n_m} \setminus \{0\})$$

with $n_i \geq 2$. The group $G = G_1 \times \ldots \times G_m$, where every component $G_i$ is either $\mathrm{SL}(n_i)$ or $\mathrm{Sp}(n_i)$, and $n_i$ is even in the second case, acts on $\mathcal{X}$ transitively and effectively. Let $\mathbb{S} = (\mathbb{K}^\times)^m$ be an algebraic torus acting on $\mathcal{X}$ by component-wise scalar multiplication, and

$$p \colon \mathcal{X} \to \mathcal{Y} := \mathbb{P}^{n_1-1} \times \cdots \times \mathbb{P}^{n_m-1}$$

be the quotient morphism. Fix a closed subgroup $S \subseteq \mathbb{S}$. The action of the group $S$ on $\mathcal{X}$ admits a geometric quotient $p_X \colon \mathcal{X} \to X := \mathcal{X}/S$. The variety $X$ is toric, it carries the induced action of the quotient group $\mathbb{S}/S$, and there is a quotient morphism $p^X \colon X \to \mathcal{Y}$ for this action closing the commutative diagram

$$\begin{tikzcd}
\mathcal{X} \arrow[rr, "p_X"] \arrow[dr, "p"'] & & X \arrow[dl, "p^X"] \\
& \mathcal{Y} &
\end{tikzcd}$$

The induced action of the group $G$ on $X$ is transitive and locally effective. We say that the $G$-variety $X$ is obtained from $\mathcal{X}$ by *central factorization*. The following theorem gives a classification of transitive actions of semisimple groups on toric varieties up to a twist by a diagram automorphism of the acting group.

---

2010 *Mathematics Subject Classification.* Primary 14M17, 14M25; Secondary 14L30.

*Key words and phrases.* Toric variety, algebraic group, homogeneous space, Cox construction.

Supported by RFBR grants 09-01-00648-a, 08-01-91855KO-a, and the Deligne 2004 Balzan prize in Mathematics.

**Theorem 1.** *Let $X$ be a toric variety with a transitive locally effective action of a connected simply connected semisimple algebraic group $G$. Then $G = G_1 \times \ldots \times G_m$, where every simple component $G_i$ is either $\mathrm{SL}(n_i)$ or $\mathrm{Sp}(n_i)$, and the variety $X$ is obtained from $\mathcal{X} = \mathcal{X}(n_1, \ldots, n_m)$ by central factorization. Conversely, any variety obtained from $\mathcal{X}$ by central factorization is a homogeneous toric variety.*

Theorem 1 also describes homogeneous spaces of a semisimple group that have a toric structure. It is natural to apply the Cox realization of a variety in order to search for toric varieties in a given class of varieties. This idea is already used in [7], where toric affine $\mathrm{SL}(2)$-embeddings are characterized.

In Section 2 we recall basic facts on the Cox realization and its generalization. Criterions of existence of an open $G$-orbit on $X$ in terms of $G$- and $(G \times S)$-actions on the total coordinate space $Z$ are also given there. In Section 3 we prove Theorem 1. The next section is devoted to special classes of toric homogeneous varieties and to a characterization of their fans. In the last section we consider transitive actions of reductive groups on toric varieties.

Our results are closely connected with the results of E.B. Vinberg [17], where algebraic transformation groups of maximal rank were classified. Recall that an *algebraic transformation group of maximal rank* is an effective generically transitive (i.e., with an open orbit) action of an algebraic group $\mathcal{G}$ on an algebraic variety $X$ such that $\dim X = \mathrm{rk}\,\mathcal{G}$, where $\mathrm{rk}\,\mathcal{G}$ is the rank of a maximal torus $T$ of the group $\mathcal{G}$. In this situation the induced action of the torus $T$ on $X$ is effective and generically transitive, see [6]. If the group $\mathcal{G}$ is semisimple, then an open $\mathcal{G}$-orbit on $X$ is a homogeneous toric variety. It turns out that in this case $X$ is a product of projective spaces and $\mathcal{G}$ acts on $X$ transitively. Theorem 1 implies that every homogeneous toric variety determines a reductive transformation group of maximal rank; here $\mathcal{G}$ is the quotient group $(\mathrm{GL}(n_1) \times \ldots \times \mathrm{GL}(n_m))/S$.

Finally, let us mention a related result from toric topology. A *torus manifold* is a smooth real even-dimensional manifold $M^{2n}$ with an effective action of a compact torus $(S^1)^n$ such that the set of $(S^1)^n$-fixed points is nonempty. In [15], homogeneous torus manifolds are studied. The latter are torus manifolds $M^{2n}$ with a transitive action of a compact Lie group $K$ such that the induced action of a maximal torus of $K$ coincides with the given $(S^1)^n$-action. It is proved that every homogeneous torus manifold may be realized as

$$M = \mathbb{CP}^{n_1} \times \ldots \times \mathbb{CP}^{n_k} \times (S^{2m_1} \times \ldots \times S^{2m_l})/F,$$

where $S^{2m}$ is a sphere of dimension $2m$, $F$ is a subgroup of $\mathbb{Z}_2 \times \ldots \times \mathbb{Z}_2$ ($l$ copies), and each copy of $\mathbb{Z}_2$ acts on the corresponding sphere by central symmetry. A compact Lie group

$$K = \mathrm{PSU}(n_1+1) \times \ldots \times \mathrm{PSU}(n_k+1) \times \mathrm{SO}(2m_1+1) \times \ldots \times \mathrm{SO}(2m_l+1)$$

acts on $M$ transitively. Moreover, the manifold $M$ is orientable if and only if $F \subset \mathrm{SO}(2m_1 + 2m_2 + \ldots + 2m_l + l)$.

The authors are grateful to E.B.Vinberg and to the referee for useful comments and suggestions.

## 2. The Cox construction

A *toric variety* is a normal algebraic variety with an effective generically transitive action of an algebraic torus $T$. A toric variety $X$ is *non-degenerate* if any invertible regular function on $X$ is constant.

Let $\mathrm{Cl}(X)$ be the divisor class group of the variety $X$. It is well-known that the group $\mathrm{Cl}(X)$ of a toric variety $X$ is finitely generated, see [11, Section 3.4]. Recall that a *quasitorus* is an affine algebraic group $S$ isomorphic to a direct product of an algebraic torus $S^0$ and a finite abelian group $\Gamma$. Every closed subgroup of a torus is a quasitorus. The group of characters of a quasitorus $S$ is a finitely generated abelian group. The *Neron-Severi quasitorus* of a toric variety $X$ is a quasitorus $S$ whose group of characters is identified with $\mathrm{Cl}(X)$.

We come to a canonical quotient realization of a non-degenerate toric variety $X$ obtained in [5]. Let $d$ be the number of prime $T$-invariant Weil divisors on $X$. Consider the vector space $\mathbb{K}^d$ and the torus $\mathbb{T} = (\mathbb{K}^\times)^d$ of all invertible diagonal matrices acting on $\mathbb{K}^d$. Then there are a closed embedding of the Neron-Severi quasitorus $S$ into $\mathbb{T}$ and an open subset $U \subseteq \mathbb{K}^d$ such that

- the complement $\mathbb{K}^d \setminus U$ is a union of some coordinate subspaces of dimension $\leq d-2$;
- there exist a categorical quotient $p_X \colon U \to U/\!\!/S$ and an isomorphism $\varphi \colon X \to U/\!\!/S$;
- via isomorphism $\varphi$, the $T$-action on $X$ corresponds to the action of the quotient group $\mathbb{T}/S$ on $U/\!\!/S$.

Later this realization was generalized to a wider class of normal algebraic varieties, see [10], [4], [9]. One of the conditions that determines this class is finite generation of the divisor class group $\mathrm{Cl}(X)$. This allows to define the Neron-Severi quasitorus $S$ of the variety $X$. The space $\mathbb{K}^d$ is replaced by an affine factorial (or, more generally, factorially graded, see [1]) $S$-variety $Z$. It is called the *total coordinate space* of the variety $X$. Further, $X$ appears as the quotient space of the categorical quotient $p_X \colon U \to U/\!\!/S$, where $U$ is an open $S$-invariant subset of $Z$ such that the complement $Z \setminus U$ is of codimension at least two in $Z$. The morphism $p_X \colon U \to X \cong U/\!\!/S$ is called the *universal torsor* over $X$.

Let a connected affine algebraic group $G$ act on a normal variety $X$. Passing to a finite covering we may assume that $\mathrm{Cl}(G) = 0$ [12, Proposition 4.6]. Then the action of $G$ on $X$ can be lifted to an action of $G$ on the total coordinate space $Z$ that commutes with the $S$-action, see [3, Section 4]. It turns out that the set $U$ is $(G \times S)$-invariant and $p_X \colon U \to X$ is a $G$-equivariant morphism.

**Lemma 1.** *The following conditions are equivalent.*

(i) *The action of the group $G$ on $X$ is generically transitive.*
(ii) *The action of the group $G \times S$ on $Z$ is generically transitive.*

*Proof.* Let $X_0 \subseteq X$ be an open $G$-orbit. Each point $x \in X_0$ is smooth on $X$, and thus the fiber $p_X^{-1}(x)$ is isomorphic to the quasitorus $S$ [9, Proposition 2.2, (iii)]. It shows that the group $G \times S$ acts on $p_X^{-1}(X_0)$ transitively.

Conversely, if $Z_0 \subseteq Z$ is an open $(G \times S)$-orbit, then $Z_0 \subseteq U$ and the action of $G$ on the quotient space $U/\!\!/S$ is generically transitive. □

Assume that the group $G$ has trivial group of characters. Then the lifting of the action of the group $G$ to $Z$ is unique, compare [3, Remark 4.1] and [8, Proposition 1.8]. Let $H$ be a closed subgroup of $G$. Every invertible regular function on the homogeneous space $G/H$ is constant, see [13, Proposition 1.2].

**Proposition 1.** *The following conditions are equivalent.*

(i) *The action of the group $G$ on $X$ is generically transitive and the complement of an open $G$-orbit has codimension at least two in $X$.*

(ii) *The action of the group $G$ on the total coordinate space $Z$ is generically transitive.*
(iii) *The action of the group $G$ on the total coordinate space $Z$ is generically transitive and the complement of an open $G$-orbit has codimension at least two in $Z$.*

*Proof.* We check "(i) $\Rightarrow$ (iii)". Let $X_0 \subseteq X$ be an open $G$-orbit. The condition $\mathrm{codim}_X(X \setminus X_0) \geq 2$ implies that $p_X\colon p_X^{-1}(X_0) \to X_0$ is the universal torsor over $X_0$ and that the complement to $p_X^{-1}(X_0)$ in $Z$ does not contain divisors, see [2, Section 2]. By [2, Lemma 3.14] (see also [1, Theorem 4.1]), the universal torsor over a homogeneous space $G/H$ is the projection $G/H_1 \to G/H$, where $H_1$ is the intersection of kernels of all characters of the subgroup $H$. This shows that the group $G$ acts on $p_X^{-1}(X_0)$ transitively.

In order to obtain "(iii) $\Rightarrow$ (i)" note that $p_X(Z_0)$, where $Z_0$ is the open $G$-orbit in $Z$, is an open $G$-orbit in $X$ whose complement does not contain divisors. The implication "(iii) $\Rightarrow$ (ii)" is obvious.

To verify "(ii) $\Rightarrow$ (iii)" let $Z_0 \subseteq Z$ be an open $G$-orbit. Since the subset $Z_0$ is $S$-invariant, for every prime divisor $D \subset Z$ in the complement to $Z_0$ the set $S \cdot D$ is an $S$-invariant Weil divisor. Each $S$-invariant Weil divisor on $Z$ is a principal divisor $\mathrm{div}(f)$ of a regular function $f \in \mathbb{K}[Z]$, see [9, Proposition 2.2, (iv)]. Then the non-constant function $f$ is invertible on $Z_0$, a contradiction. □

The same arguments lead to the following result.

**Proposition 2.** *The action of the group $G$ on $X$ is transitive if and only if the open subset $U \subseteq Z$ is a $G$-orbit.*

## 3. CLASSIFICATION OF HOMOGENEOUS TORIC VARIETIES

In this section we prove Theorem 1. Since the variety $X$ is toric, its total coordinate space $Z$ is an affine space.

**Lemma 2.** *Let a semisimple group $G$ act on a toric variety $X$ with an open orbit. Then $X$ is non-degenerate and the action of the group $G \times S$ on the affine space $Z$ is equivalent to a linear one.*

*Proof.* Since any invertible function of the open $G$-orbit is constant, the variety $X$ is non-degenerate. By Lemma 1, the action of the group $G \times S$ on the space $Z$ is generically transitive, and the second statement follows from [14, Proposition 5.1]. □

Later on we assume that $G = G_1 \times \ldots \times G_m$ acts on $X$ transitively. Denote by $V$ the total coordinate space $Z$ of the variety $X$ regarded as the $(G \times S)$-module. We proceed with a description of the $G$-module structure on $V$.

**Proposition 3.** *Let $V = V_1 \oplus \ldots \oplus V_s$ be a decomposition into irreducible summands. Then every simple component $G_i$ acts not identically only on one summand $V_i$ (up to renumbering), and thus $m = s$. Moreover, every $G_i$ acts on the set of nonzero vectors in $V_i$ transitively.*

*Proof.* By Proposition 2, the complement of the open $G$-orbit $U$ in $V$ is a union of coordinate subspaces (in some, possibly nonlinear, coordinate system). Thus each irreducible component of the complement is a smooth variety. The linear action of the group $G$ on $V$ commutes with the group $\mathbb{K}^\times$ of scalar operators, and the open orbit $U$ as well as any component of the complement $V \setminus U$ is $(G \times \mathbb{K}^\times)$-invariant. But a cone is a smooth variety if and only if it is a subspace. This shows that each component of $V \setminus U$ is a maximal proper submodule of $V$. In particular, the number of maximal proper submodules is finite and thus the $G$-modules $V_1, \ldots, V_s$ are pairwise non-isomorphic. The orbit $U$ is the set of vectors $v \in V$ whose projection on each $V_i$ is nonzero. This implies that the group $G$ acts on the set of nonzero vectors of each submodule $V_i$ transitively.

If several components of $G$ act on some $V_i$ not identically, then $V_i$ is isomorphic to the tensor product of simple modules of these components. Then the cone of decomposable tensors in $V_i$ is $G$-invariant, a contradiction.

Suppose that a simple component $G_l$ acts on both $V_i$ and $V_j$ not identically. Then $G_l$ acts transitively on the set of pairs $(v_i, v_j)$ with nonzero $v_i$ and $v_j$. In particular, any such pair is an eigenvector of a Borel subgroup of $G_l$. Fix a Borel subgroup $B \subset G_l$ and a highest vector for $B$ in $V_i$ as $v_i$ and a lowest vector for $B$ in $V_j$ as $v_j$. Since the intersection of two opposite parabolic subgroups of $G_l$ does not contain a Borel subgroup, we get a contradiction. ☐

The following lemma is well known. We give a short self-contained proof suggested by the referee.

**Lemma 3.** *Finite-dimensional rational modules of a simple group $G$ such that $G$ acts on the set of nonzero vectors transitively are*

(1) *the tautological* $\mathrm{SL}(n)$*-module* $\mathbb{K}^n$ *and* $\mathrm{Sp}(2n)$*-module* $\mathbb{K}^{2n}$*;*
(2) *the dual* $\mathrm{SL}(n)$*-module* $(\mathbb{K}^n)^*$*.*

*Proof.* Since $G$ acts on $V \setminus \{0\}$ transitively, $V$ is a simple $G$-module of highest weight $\lambda$ and $V = \mathfrak{g}v_\lambda$, where $\mathfrak{g}$ is the tangent algebra of the group $G$ and $v_\lambda$ is a highest weight vector. In particular, a lowest weight vector is $v_{-\lambda^*} = e_{-\alpha}v_\lambda$, where $\alpha$ is a positive root, whence $\alpha = \lambda + \lambda^*$ is the highest root. This occurs only for $G = \mathrm{SL}(n)$ with fundamental weights $\lambda = \omega_1, \omega_{n-1}$, and $G = \mathrm{Sp}(2n)$ with $\lambda = \omega_1$. ☐

Applying an outer automorphism of $G$, we may assume that $G = G_1 \times \ldots \times G_m$ and $V = V_1 \oplus \ldots \oplus V_m$, where every component $G_i$ is either $\mathrm{SL}(n_i)$ or $\mathrm{Sp}(n_i)$, and $V_i$ is the tautological $G_i$-module with identical action of other components. The open $G$-orbit $U$ in $V$ coincides with the subvariety $\mathcal{X} = \mathcal{X}(n_1, \ldots, n_m)$. Therefore the variety $X$ is obtained from $\mathcal{X}$ by central factorization.

Let $\mathbb{S} = (\mathbb{K}^\times)^m$ be an algebraic torus acting on $V = V_1 \oplus \ldots \oplus V_m$ by component-wise scalar multiplication. It remains to explain why for any subgroup $S \subseteq \mathbb{S}$ there exists a geometric quotient $\mathcal{X} \to \mathcal{X}/S$. This follows from the fact that $\mathcal{X}$ is a homogeneous space of the group $\overline{G} := \mathrm{GL}(n_1) \times \ldots \times \mathrm{GL}(n_m)$, and $S$ is a central subgroup of $\overline{G}$. The proof of Theorem 1 is completed.

*Remark* 1. The collection $(n_1, \ldots, n_m)$ is determined by a homogeneous toric variety $X$ uniquely. Indeed, if $\mathbb{K}^d \supset U \to X$ is the Cox realization of $X$ and $C_1, \ldots, C_m$ are irreducible components of the complement $\mathbb{K}^d \setminus U$, then $n_i = d - \dim C_i$.

## 4. Properties of homogeneous toric varieties

In this section we use standard notation of toric geometry, see [11]. Let $\mathcal{N}$ be the lattice of one-parameter subgroups of a $d$-dimensional torus $\mathbb{T}$ and $\mathcal{M}$ be the lattice of characters of $\mathbb{T}$. The torus $\mathbb{T}$ acts diagonally on the space $\mathbb{K}^d = V = V_1 \oplus \ldots \oplus V_m$, and $\mathbb{S} \subset \mathbb{T}$ is the

$m$-dimensional subtorus acting on every $V_i$ by scalar multiplication. Identification of $\mathbb{T}$ with $(\mathbb{K}^\times)^d$ defines standard bases in $\mathcal{N}$ and $\mathcal{M}$. Moreover, the decomposition $V = V_1 \oplus \ldots \oplus V_m$ divides the standard basis of $\mathcal{N}$ into $m$ groups $I_1, \ldots, I_m$, where each group $I_j$ contains $n_j$ basis vectors and $n_j := \dim V_j$. The open subvariety $\mathcal{X}(n_1, \ldots, n_m) = U \subset V$ is a toric $\mathbb{T}$-variety. Its fan $\mathcal{C} = \mathcal{C}(n_1, \ldots, n_m)$ in the lattice $\mathcal{N}$ consists of the cones generated by all collections of standard basis vectors that do not contain any subset $I_j$.

Let $S \subseteq \mathbb{S}$ be a closed subgroup. There is a sequence of lattices of one-parameter subgroups $\mathcal{N}_S \subseteq \mathcal{N}_{\mathbb{S}} \subset \mathcal{N}$, where the lattice $\mathcal{N}_S$ is determined by the connected component $S^0$ of the quasitorus $S$. The fan $\mathcal{C}_{S^0}$ of the quotient space $\mathcal{X}/S^0$ is the image of the fan $\mathcal{C}$ under the projection

$$\mathcal{N}_{\mathbb{Q}} \ \to \ (\mathcal{N}/\mathcal{N}_S)_{\mathbb{Q}}.$$

The fan $\mathcal{C}_S$ of the variety $\mathcal{X}/S$ coincides with the fan $\mathcal{C}_{S^0}$ considered with regard to an overlattice of $\mathcal{N}/\mathcal{N}_S$ of finite index, see [11, Section 2.2]. In particular, the fan $\mathcal{C}_{\mathbb{S}}$ coincides with the fan $\mathcal{P}$ of the product of projective spaces $\mathbb{P}^{n_1-1} \times \ldots \times \mathbb{P}^{n_m-1}$, and $\mathcal{C}_S$ may be considered as an intermediate step of the projection:

$$\mathcal{C} \to \mathcal{C}_S \to \mathcal{P}.$$

Let us define a sublattice $\mathcal{M}_S \subseteq \mathcal{M}$ as the set of characters of the torus $\mathbb{T}$ containing $S$ in the kernel. Elements of $\mathcal{M}_S$ are linear functions on the space $(\mathcal{N}/\mathcal{N}_S)_{\mathbb{Q}}$.

**Proposition 4.** *Let $X = \mathcal{X}/S$ be a homogeneous toric variety. Then*

1. *the variety $X$ is quasiprojective;*
2. *the variety $X$ is not affine;*
3. *the variety $X$ is projective if and only if it coincides with $\mathbb{P}^{n_1-1} \times \ldots \times \mathbb{P}^{n_m-1}$;*
4. *the variety $X$ is quasiaffine if and only if the lattice $\mathcal{M}_S$ contains a vector with positive coordinates;*
5. *the variety $X$ has a nonconstant regular function if and only if the lattice $\mathcal{M}_S$ contains a nonzero vector with nonnegative coordinates.*

*Proof.* (1) By Chevalley's Theorem, any homogeneous space of an affine algebraic group is a quasiprojective variety.

(2) A toric variety obtained via Cox construction is affine if and only if $U = V$. In our situation this is not the case.

(3) Maximal dimension of a cone in the fan $\mathcal{C}$ equals $n_1 + \ldots + n_m - m$. Therefore the fan $\mathcal{C}_S$ is complete if and only if it is obtained from $\mathcal{C}$ by projection to $(\mathcal{N}/\mathcal{N}_{\mathbb{S}})_{\mathbb{Q}}$, and thus $\mathcal{C}_S$ coincides with $\mathcal{P}$.

(4) A toric variety is quasiaffine if and only if its fan is a collection of faces of a strongly convex polyhedral cone. In our case, this condition implies that the projection $K$ of the support of the fan $\mathcal{C}$ to $(\mathcal{N}/\mathcal{N}_S)_{\mathbb{Q}}$ is a strongly convex cone. The latter is equivalent to existence of a linear function on the space $(\mathcal{N}/\mathcal{N}_S)_{\mathbb{Q}}$ that is positive on $K \setminus \{0\}$. This gives the desired element of the lattice $\mathcal{M}_S$.

Conversely, assume that the lattice $\mathcal{M}_S$ contains a vector $v$ with positive coordinates. We have to show that the projection of each cone of the fan $\mathcal{C}$ is a face of $K$. Fix proper subsets $J_1 \subset I_1, \ldots, J_m \subset I_m$ of the sets of standard basis vectors of the lattice $\mathcal{N}$. We claim that there is an element of the lattice $\mathcal{M}_S$, which vanishes on the vectors of $J_1 \cup \ldots \cup J_m$ and is positive on other standard basis vectors. Indeed, the sublattice $\mathcal{M}_S$ is defined in terms of the sums of coordinates of a character over all $m$ groups of its coordinates. The desired vector should have the same sums of coordinates over the groups as the vector $v$.

(5) Since regular functions on $X$ form a rational $\mathbb{T}$-module, one may consider only $\mathbb{T}$-semiinvariant regular functions. Further, regular $\mathbb{T}$-semiinvariants on $X$ correspond to characters from $\mathcal{M}_S$ that are nonnegative on the rays of the fan $\mathcal{C}$, see [11, Section 3.3]. □

*Remark* 2. Let $X$ be a homogeneous toric variety. Then $X$ is projective if and only if $X$ contains a $\mathbb{T}$-fixed point. Indeed, the latter condition means that the fan $\mathcal{C}_S$ contains a cone of full dimension, thus $\mathcal{N}_S = \mathcal{N}_{\mathbb{S}}$ and $S = \mathbb{S}$.

**Example 1.** Let $m = 2$ and $n_1 = n_2 = 2$. Then $\mathcal{X} = (\mathbb{K}^2 \setminus \{0\}) \times (\mathbb{K}^2 \setminus \{0\})$. Set $S = \{(s,s,s,s) : s \in \mathbb{K}^\times\}$. Then

$$\mathcal{M}_S = \{(x_1,x_2,x_3,x_4)\,;\, x_i \in \mathbb{Z},\ x_1+x_2+x_3+x_4 = 0\},$$

and the variety $X$ is $\mathbb{P}^3 \setminus (D_1 \cup D_2)$, where $D_i \cong \mathbb{P}^1$. If we set $S = \{(s,s,s^{-1},s^{-1}) : s \in \mathbb{K}^\times\}$, then

$$\mathcal{M}_S = \{(x_1,x_2,x_3,x_4)\,;\, x_i \in \mathbb{Z},\ x_1+x_2 = x_3+x_4\},$$

and $X$ is a three-dimensional quadratic cone with the apex removed.

Let us characterize the fans of homogeneous toric varieties. Let $N$ be a lattice, $\Delta$ be a fan in $N_{\mathbb{Q}}$ and $P$ be the set of primitive vectors on the rays of $\Delta$. Denote by $N_0$ a sublattice of $N$ generated by $P$. Fix a positive integer $m$.

**Definition 1.** A fan $\Delta$ is called *$m$-partite* if

- the set $P$ spans the vector space $N_{\mathbb{Q}}$;
- the set $P$ can be decomposed into $m$ subsets $P = I_1 \sqcup \ldots \sqcup I_m$, where each $I_j$ contains at least two elements, and the cones of $\Delta$ are exactly the cones generated by subsets $J \subset P$ that do not contain any $I_j$.

Set $I_j = \{e^j_1, \ldots, e^j_{n_j}\}$ and $q_j = e^j_1 + \ldots + e^j_{n_j}$. Let $Q$ be a sublattice of $N$ generated by $q_1, \ldots, q_m$, and $Q_{\mathbb{Q}} = Q \otimes_{\mathbb{Z}} \mathbb{Q}$.

**Proposition 5.** *A fan $\Delta$ is the fan of a homogeneous toric variety if and only if*

1. *$\Delta$ is $m$-partite for some $m \geq 1$;*
2. *every linear relation among elements of $P$ has the form $\lambda_1 q_1 + \ldots + \lambda_m q_m = 0$ for some rational $\lambda_i$;*
3. *$N \subset N_0 + Q_{\mathbb{Q}}$.*

*Proof.* A fan is $m$-partite if and only if it is a projection of the fan $\mathcal{C}(n_1, \ldots, n_m)$ with some $n_i \geq 2$. Condition 2 means that the kernel of the projection is of the form $(\mathcal{N}_{S^0})_{\mathbb{Q}}$, where $S \subseteq \mathbb{S}$. Finally, condition 3 means that $N$ is generated by $P$ and some elements

$$\frac{r_{1i}}{R_i} q_1 + \ldots + \frac{r_{mi}}{R_i} q_m, \quad \text{where } r_{ji} \in \mathbb{Z}_{\geq 0},\ \ R_i \in \mathbb{Z}_{>0},\ \ r_{ji} < R_i, \ \text{ and } \ i = 1, \ldots, l.$$

Equivalently, the corresponding toric variety is obtained as the quotient of the variety $\mathcal{X}(n_1, \ldots, n_m)/S^0$ by an action of the group $\Gamma = \Gamma_1 \times \ldots \times \Gamma_l$, where $\Gamma_i$ is the cyclic group of $R_i$-th roots of unity and an element $\epsilon \in \Gamma_i$ multiplies the $j$-th factor of $\mathcal{X}(n_1, \ldots, n_m)$ by $\epsilon^{r_{ji}}$. □

## 5. Some generalizations

Let a connected reductive group $G$ act on a toric variety $X$ transitively. One may assume that $G = G^s \times L$, where $G^s$ is a simply connected semisimple group, $L$ is a central torus, and the $G$-action on $X$ is locally effective. It is well known that any toric variety $X$ is isomorphic to a direct product $X_0 \times X_1$, where $X_0$ is a non-degenerate toric variety and $X_1$ is an algebraic torus.

Let us give a construction of a transitive $G$-action on a toric variety $X$. Take a $G^s$-homogeneous toric variety $X_0$ with a locally effective and $G^s$-equivariant action of a quasitorus $L'$. Fix an inclusion $L' \subseteq L$ into an algebraic torus $L$ as a closed subgroup. The group $G = G^s \times L$ acts on $X_0 \times L$, where $G^s$ acts on the first factor and $L$ acts on the second one by multiplication. Consider the $G$-equivariant action of $L'$ on $X_0 \times L$ given by $(x_0, l) \mapsto (sx_0, s^{-1}l)$ for every $s \in L'$. Then

$$X(X_0, G^s, L', L) := (X_0 \times L)/L'$$

is a $G$-homogeneous toric variety.

**Proposition 6.** *Let $X$ be a toric variety endowed with a transitive and locally effective action of a connected reductive group $G = G^s \times L$. Then the non-degenerate factor $X_0$ of $X$ is a $G^s$-homogeneous toric variety. Moreover, if $L'$ is the stabilizer of a $G^s$-orbit on $X$ in the torus $L$, then $X$ is $G$-equivariantly isomorphic to $X(X_0, G^s, L', L)$.*

*Proof.* Since the $G^s$- and $L$-actions on $X$ commute, all $G^s$-orbits are of the same dimension. Let $Y$ be one of these orbits. Any invertible function on $Y$ is constant. Consider the above decomposition $X = X_0 \times X_1$. Since points on $X_1$ are separated by invertible functions, $Y$ is contained in a subvariety $X_0 \times \{x_1\}$, where $x_1 \in X_1$. Let $L'$ be the stabilizer of the subvariety $Y$ in the torus $L$. Then the stabilizer $H$ of a point $x \in Y$ is contained in the subgroup $G^s \times L'$ and the homogeneous space $G/H$ projects onto $G/(G^s \times L') \cong L/L'$. Points on $L/L'$ are separated by invertible functions, hence $X_0 \times \{x_1\}$ is contained in a fiber of the projection. But the fibers coincide with $G^s$-orbits on $X$. This implies $Y = X_0 \times \{x_1\}$.

Let us identify the variety $X_0$ with the subvariety $Y \subseteq X$. Consider the morphism

$$\varphi \colon X_0 \times L \to X, \quad (x_0, l) \mapsto lx_0.$$

Two pairs $(x_0, l)$ and $(\widetilde{x_0}, \widetilde{l})$ are in the same fiber of $\varphi$ if and only if $(\widetilde{x_0}, \widetilde{l}) = (sx_0, s^{-1}l)$ with $s = \widetilde{l}^{-1}l$. This shows that $\varphi$ induces a bijective morphism $X(X_0, G^s, L', L) \to X$. Clearly, this is an isomorphism of $G$-homogeneous spaces. $\square$

If the subgroup $L'$ is connected, then $L \cong L' \times L''$ with some complementary subtorus $L''$, and $X \cong X_0 \times L''$. But unlike the case of algebraic transformation groups of maximal rank [17, Theorem 2], this situation does not always occur. Indeed, one may consider a toric variety $(\mathbb{K}^2 \setminus \{0\}) \times \mathbb{K}^\times$ with a transitive locally effective action of the group $\mathrm{SL}(2) \times \mathbb{K}^\times$ given as $(g, t) \cdot (v, a) = (g(tv), t^2 a)$.

*Remark* 3. It would be interesting to generalize [17, Theorem 3] and to describe toric varieties with transitive actions of non-reductive affine algebraic groups.

Besides homogeneous toric varieties, our method allows to describe toric varieties with a generically transitive action of a semisimple group $G$. By Lemma 1, they are quasitorus quotients of open subsets of generically transitive $(G \times \mathbb{S})$-modules. Such modules are known as $(G \times \mathbb{S})$-prehomogeneous vector spaces. For an explicit description, one needs a list of prehomogeneous vector spaces. The classification results here are known only under some restrictions on the group and on the module. For example, if $G$ is simple and the number of irreducible summands of the module does not exceed three, the classification is given in a series of papers of M.Sato, T.Kimura, K.Ueda, T.Yoshigiaki and others.

If the complement of an open $G$-orbit on a toric variety $X$ has codimension at least two in $X$, then $X$ comes from a $G$-prehomogeneous vector space (Proposition 1). When the group $G$ is simple, the list of $G$-prehomogeneous vector spaces is obtained in [16, Theorems 7-8], and the corresponding toric varieties are described in [2, Proposition 4.7]. In constrast to the homogeneous case, here appear singular [2, Example 5.8] and non-quasiprojective [2, Example 5.9] varieties.

## References

[1] Arzhantsev, I.V., *On the factoriality of Cox rings*, Mat. Zametki **85** (2009), 643–651 (Russian); English transl.: Math. Notes **85** (2009), 623–629.

[2] Arzhantsev, I.V., and J. Hausen, *On embeddings of homogeneous spaces with small boundary*, J. Algebra **304** (2006), 950–988.

[3] Arzhantsev, I.V., and J. Hausen, *Geometric Invariant Theory via Cox rings*, J. Pure Appl. Algebra **213** (2009), 154–172.

[4] Berchtold, F., and J. Hausen, *Cox rings and combinatorics*, Trans. AMS **359** (2007), 1205–1252.

[5] Cox, D.A., *The homogeneous coordinate ring of a toric variety*, J. Alg. Geometry **4** (1995), 17–50.

[6] Demazure, M., *Sous-groupes algébriques de rang maximum du groupe de Cremona*, Ann. Sci. École Norm. Sup. (4) **3** (1970), 507–588.

[7] Gaifullin, S.A., *Affine toric* SL(2)*-embeddings*, Mat. Sbornik **199** (2008), 3–24 (Russian); English transl.: Sbornik Math. **199** (2008), 319–339.

[8] Hausen, J., *Geometric invariant theory based on Weil divisors*, Compos. Math. **140** (2004), 1518–1536.

[9] Hausen, J., *Cox rings and combinatorics, II*, Moscow Math. J. **8** (2008), 711–757.

[10] Hu, Y., and S. Keel, *Mori dream spaces and GIT*, Michigan Math. J. **48** (2000), 331–348.

[11] Fulton, W., Introduction to toric varieties, Annals of Math. Studies 131, Princeton University Press, Princeton, NJ, 1993.

[12] Knop, F., H. Kraft, D. Luna, and Th. Vust, *Local properties of algebraic group actions*, Algebraische Transformationsgruppen und Invariantentheorie, DMV Sem., **13**, Birkhäuser, Basel (1989), 63–75.

[13] Knop, F., H. Kraft, and Th. Vust, *The Picard group of a G-variety*, Algebraische Transformationsgruppen und Invariantentheorie, DMV Sem., **13**, Birkhäuser, Basel (1989), 77–87.

[14] Kraft, H., and V.L. Popov, *Semisimple group actions on the three-dimensional affine space are linear*, Comment. Math. Helv. **60** (1985), 466–479.

[15] Kuroki, Sh., *Classification of homogeneous torus manifolds*, Preprint, Fudan University, 2007.

[16] Vinberg, E.B., *Invariant linear connections in a homogeneous space*, Trudy Moskov. Mat. Obsch. **9** (1960), 191–210 (Russian).

[17] Vinberg, E.B., *Algebraic transformation groups of maximal rank*, Mat. Sbornik (N.S.) **88:130** (1972), 493–503 (Russian).

Ivan V. Arzhantsev, Department of Higher Algebra, Faculty of Mechanics and Mathematics, Moscow State University, Leninskie Gory 1, Moscow, 119991, Russia
*E-mail address*: `arjantse@mccme.ru`

Sergey A. Gaifullin, Department of Higher Algebra, Faculty of Mechanics and Mathematics, Moscow State University, Leninskie Gory 1, Moscow, 119991, Russia
*E-mail address*: `sgayf@yandex.ru`